

Surprising competition found in high-temperature superconductors

A team led by SLAC National Accelerator Laboratory and Stanford University scientists has made an important discovery toward understanding how a large group of complex copper oxide materials lose their electrical resistance at remarkably high temperatures.

The materials in question are high-temperature superconductors, which conduct electricity perfectly with no resistance when cooled below -100 C.

In a report published in the *Proceedings of the National Academy of Sciences (PNAS)*, researchers led by SLAC Chief Scientist and Stanford Professor Zhi-Xun Shen describe the surprisingly complex and dynamic way that the electrons organize themselves within one group of copper oxide superconductors, known as cuprates. They created a complete phase diagram of the material—a roadmap of its properties—over a range of compositions and temperatures that are ideal for superconductivity.

Earlier experiments had shown that non-superconducting behavior—the so-called pseudogap phase—can occur within cuprate superconductors. In this study, the researchers were surprised to discover that this pseudogap phase actually coexists and competes with the superconducting phase over a wide range of temperatures and compositions.

"The pseudogap is one of the biggest mysteries in high-temperature superconductivity," says Inna Vishik, a Stanford graduate student in Shen's group who is first author on the *PNAS* paper. "We had thought that the pseudogap did not notice the onset of superconductivity. But that is not the case. As temperature decreases, superconductivity actually suppresses the pseudogap phase."

Learning the details of this competition between the pseudogap and superconducting phases should aid researchers in their ultimate goal of creating new superconducting materials that operate above room temperature—an advancement that would revolutionize myriad modern technologies, from tiny computer chips to miles-long power-transmission lines.

"Twenty-five years after the discovery of high-temperature superconductivity in cuprates, this experiment has brought out important new aspects of the phase diagram that had been hidden," Shen says. "Since the phase diagram is fundamental to understanding matter, describing the nature of the phases and the intricate relationship among them, this work is an important step forward."

The complex electronic structure of cuprate superconductors has made it very difficult for scientists to determine exactly how these materials lose their electrical

Surprising competition found in high-temperature superconductors

Published on Research & Development (<http://www.rdmag.com>)

resistance, let alone how to modify them to make this transition occur above room temperature.

In recent years, Shen's research group has helped develop a technique called angle-resolved photoemission spectroscopy (ARPES), which enables scientists to see electron interactions in much more detail than before.

To explore the properties of the pseudogap and superconducting states, Vishik systematically varied the composition of a superconducting cuprate containing bismuth, strontium and calcium and performed hundreds of ARPES measurements on those samples in Shen's Stanford laboratory and at SLAC's Stanford Synchrotron Radiation Lightsource.

Future research is aimed at nailing down the exact electron behavior of the pseudogap phase and the details of its competition with the superconducting phase, as well as determining exactly how electrons pair up to create superconductivity in the copper oxides.

Source: [SLAC National Accelerator Laboratory](#) [1]

Source URL (retrieved on 05/30/2016 - 6:09pm):

<http://www.rdmag.com/news/2012/11/surprising-competition-found-high-temperature-superconductors>

Links:

[1] <http://www6.slac.stanford.edu/news/2012-11-14-high-temp-super.aspx>